


Dental Emergencies



What should you do?

A basic Guide to Dental First-Aid

Toothache - Rinse mouth with warm water. If swelling occurs, apply cold packs to outside of face. **DO NOT** apply aspirin or other medications directly to aching tooth. Go to a dentist.

Bleeding Gums - If gums are red, swollen or sore, rinse every two hours with a mixture of equal parts of water and hydrogen peroxide or with a warm salt water solution. If condition does not improve in a few days, see a dentist. Bleeding gums are often a sign of an unhealthy mouth. Be sure to brush and floss twice a day and visit a dentist every six months for a check-up.

Broken Or Dislocated Tooth - Clean injured area with warm water. If injured area begins to swell, apply cold packs to outside of face. See a dentist **immediately!**

Tongue Or Lip Bite - If bleeding, apply pressure with clean cloth. If swelling occurs, apply cold pack. If bleeding does not stop or if bite is severe, go to a hospital emergency room.

Broken Jaw - If you think the jaw is broken, **DO NOT** move the jaw. Wrap a scarf, handkerchief, tie, or towel around the head to keep the jaw from moving and go to a hospital emergency room **immediately!**

Knock Out Tooth - First, find the tooth. Place it in milk or water. Gently rinse the tooth but **DO NOT** scrub it! **DO NOT** wrap tooth in tissue! Go to the dentist with the tooth **immediately!**

Canker Or Cold Sores and Fever Blisters - For temporary pain relief, apply an over-the-counter cream/gel that has benzocaine. Benzocaine numbs the area. Follow directions on label. Avoid hot, spicy foods. A non-aspirin pill can be swallowed. **DO NOT** apply pill directly to sores. If sores do not heal within 7-14 days and/or signs of fever and pain persist, contact a dentist.

Pain From A New Tooth Appearing - Find the site and apply a cold pack to the outside of the face. A non-aspirin over-the-counter children's pain reliever can be swallowed (pill or liquid form). **DO NOT** apply pill directly on gums. An over-the-counter cream/gel that has benzocaine can also be used directly on the gums. Follow directions on medicine label. If pain continues, contact a dentist.

Objects Wedged Between Teeth - Carefully remove object from between teeth with dental floss. **DO NOT** remove sharp objects. If unable to remove or object is sharp, see a dentist **immediately!**

**If a dental emergency occurs, act fast!
Contact a dentist immediately!**



Revision 5/03



County of San Diego
Health and Human Services Agency



Brought to you by the U.S. Department of Agriculture, and the California State Department of Health Services Cancer Prevention and Nutrition Section